

Cedar River and Bear Creek Chinook

Abundance and Survival-to-Emigration
into Lake Washington of the 2000 Brood

Dave Seiler and
Greg Volkhardt

Wild Stock Production Evaluation Unit

Chinook Escapement

Cedar River – AUC Method

Year	Esc.
2001:	810
2000:	120
1999:	241
1998:	432


Contributors

Steve Foley – WDFW
King County
Muckleshoot Tribe
City of Seattle


Bear Creek – AUC Method

2001:	459	▶	690
2000:	228	▶	332
1999:	537	▶	732
1998:	265	▶	398

Juvenile Chinook Production Monitoring



Location Map of the Cedar River and Bear Creek Trap Sites



Cedar River



Wild Stock Production Evaluation Unit


Big Bear Creek



Wild Stock Production Evaluation Unit


Incubation Success

- It's a function of peak flow -




Incubation Success

- It's a function of peak flow -




Incubation Timing

2001 Cedar River Chinook 0+



Incubation Timing


2001 Bear Creek Chinook 0+






Early Rearing and Migration

Bi-modal migration timing

- ➡ "Fry" migration from January to mid-April
- ➡ "Smolt" migration from mid-April to July
- ➡ Different proportions between years



Cedar River Wild Chinook






1999 – High flows/velocities push most of the production downstream as “fry”. Low “smolt” production.

2000 – Moderate-high flows/velocities results in higher “smolt” production than observed in 1999.

2001 – Extreme low flows result in the largest “smolt” production measured. Low escapement (120) and predation contributed to the low number of total migrants.

Bear Creek Wild Chinook



1999 – Lower stream energy in Bear Creek results in high proportion of smolts even with high flow levels.

2000 – Good escapement (732) results in a higher proportion of fry being displaced downstream.

2001 – Factors such as low flow and stream energy providing an advantage to predators as well as high sockeye spawner abundance resulted in the lowest total production measured.

Cedar & Bear Chinook - Estimated production, timing, and survival to fry and smolts stages


Cedar River											
Smolt Year (I)	Estimated Migration			Percent Migration		Est Female (I-1)	Production/Female		Survival Ratios		
	Fry	Smolts	Total	Fry	Smolts		Fry	Smolts	Fry/ PED	Smolts/ PED	Total
1999	67,336	12,454	79,790	84%	16%	232	290	54	7.3%	1.3%	8.6%
2000	54,836	19,860	74,696	73%	27%	180	305	110	7.6%	2.8%	10.4%
2001	9,427	20,200	29,627	32%	68%	53	178	381	4.5%	9.5%	14.0%
Bear Creek											
Smolt Year (I)	Estimated Migration			Percent Migration		Est Female (I-1)	Production/Female		Survival Ratios		
	Fry	Smolts	Total	Fry	Smolts		Fry	Smolts	Fry/ PED	Smolts/ PED	Total
1999	1,720	13,207	14,927	12%	88%	159	11	83	0.3%	2.1%	2.4%
2000	14,116	17,785	31,901	44%	56%	293	48	61	1.2%	1.5%	2.7%
2001	541	10,616	11,157	5%	95%	133	4	80	0.1%	2.0%	2.1%

PIT Tagging Studies



Wild Stock Production Evaluation Unit

Survival through the Lake



Survival through the Lake

